

Western Burrowing Owl Workshop



Dr. Lynne Trulio
ltrulio@earthlink.net

July 19-20, 2018

Workshop Topics

- Distribution
- Identification
- Life History
- Habitat Requirements
- Status & Threats
- Habitat Enhancement Methods
- Reestablishing Owls on Sites
- Management for Population Persistence

Athene cunicularia
Burrowing Owl or "Little Miner"



An Odd Bird

- Does not hoot
- Active day and night
- Only owl that lives and nests underground
- Life revolves around the burrow
- Lines burrows with dung, collects burrow decorations
- Juveniles do a great rattlesnake mimic!



Entire Species Range - ~17 subspecies




US Subspecies of Burrowing Owls

- Two subspecies in the US:
 - Western burrowing owl (*A. cunicularia hypugaea*)
 - Florida burrowing owl (*A. cunicularia floridana*)
- Very similar in appearance/behavior



Martin Meyers

Range of the Western Burrowing Owl




Migration

- Year-round residents
- Migrants - coastal, bay edges, hill sides
- Breed in more interior, flat areas
- Focus is on the breeding season...
- ...but the winter season is also vital

RANGE MAP

Burrowing Owl



Identification - Adults

- Ht ~7.5-9.5 inches
- Wt ~5-6 ounces
- Wing span ~22 inches
- Long legs, few feathers
- Mottled brown and cream
- Designed for camouflage!
- Bright lemon yellow eyes
- No ear tufts
- Active day and night
- On ground or low perch



Identification - Adults

Male and female not sexually dimorphic, but male is slightly larger & paler in summer; behavioral differences.



Identification - Chicks

- May - September in CA
- Smaller than adults until about July
- Buffy breast, whiter eye-brow, darker collar



Steve Thurman



Chicks over the Season



Calls



- No typical owl "hoots"
- Males: "coo - coo" territorial/mating call
- All birds: "chatter" alarm call
- Females & Juveniles: "rasp" food call
- Nestlings & Juveniles: defensive call
- Listen to the calls:

http://www.allaboutbirds.org/guide/burrowing_owl/sounds

Life History Characteristics

- Inhabits open grasslands; short scrub habitat
- A raptor - although a small one
- Many predators
- Migratory in much of range, but in temperate areas some resident and some migrant
- Semi-colonial, esp. with sciurids
- Semi-fossorial - inhabits burrows year round
- Monogamous during the breeding season
- Sexually mature at 1 year
- Lays 2-12 eggs; one clutch per year
- Lives ~3-5 years, but up to ~8 years

Bird of Open Grasslands:

Prairies, Ag Lands, Bases, Golf Courses, Open Fields - Natural Grasslands and Urban Sites



Habitat Types - Statewide in CA

(Wilkerson & Siegel, 2010)

~30%=irrigation canals

~16%=natural grassland

~10%=idle/fallow field

~10%=field crop

~10%=urban

~ 8%=pasture


~ 6%=brushland

~ 3%=grain/row



But in Nevada, for example...

- 44% in sagebrush
- 22% in grasslands
- 21% in salt desert scrub
- 9% in agriculture



(Great Basin Bird Observatory.
2010. *Nevada Comprehensive
Bird Conservation Plan* at
http://www.gbbo.org/bird_conservation_plan.html)

Nesting Habitat Requirements

Flexible requirements...within limits



Nesting owls are found...

- At lower elevations in much of California (often <200 ft)
- In open areas, typically with few trees
- Short grass (<6") around burrows
- Structural heterogeneity elsewhere - long grass, shrubs, rock + brush piles
- Associated with ground squirrels
- Some level of soil disturbance, esp. from ground squirrels


Predators? Just about everything!

- Primary: hawks, larger owls, skunks, foxes, coyotes, snakes
- Others: crows, dogs, cats, badgers



Scott Thurman

Burrows are key, natural and artificial



California Ground Squirrels





Artificial
burrow with
lots of debris
out front

Many burrows are needed:

- * Per pair: primary + satellite
- * Overall: prefer high-burrow density areas



Breeding Season Territory & Home Range

- Site tenacity - during season
- Some site fidelity - 32% - 57%
- 80% of foraging within 600m of burrow, but as far as 2 miles away
- Home range size varies widely, depending on prey availability and quality


Wintering Season in CA: Many resident birds but...



- Saskatchewan Study using geolocators showed:
 - 9/10 females to CA
 - 10/15 males to OR/WA
- CA = important winter habitat for birds
- Males nearer breeding sites than females

Current
research using
satellite
telemetry
conducted by a
number of
researchers
(Conroy,
Johnston,
Holroyd,
Trefry) ...

Burrowing Owl Migration - Fall 2014





...shows amazing
migratory
travels!

Bend, Oregon


to

Salinas, California



Wintering BUOWs in the Santa Clara Valley Habitat Plan Area

- Where do BUOWs winter & how many?
- What is the relationship between wintering and breeding owls?



Santa Clara Valley Habitat Plan (HCP/NCCP)

Burrowing owls - a covered species
Seeking ways to protect & recover




SANTA CLARA VALLEY
HABITAT AGENCY




Santa Clara Valley Habitat Plan Area

--focus on protected open space--



Long-term (typical) breeding sites



Winter Study Methods

- CBC locations, local experts, eBird
- Bow trap and MP3 player
- Capture and band




Summer Study Methods


- Surveys and banding at:
 - a) Typical breeding sites and
 - b) Foothill locations of wintering owls



Winter 2014-15, 2015-16 & 2016-17


- Overall:
 - 28, 23, & 26 newly banded
 - 12-20 summer resights
 - Up to 700m in elevation
- Winter returns:
 - 2015-16: 3 from previous winter
 - 2016-17: 2 from previous winters
- Resident birds:
 - At typical breeding sites
 - No previously banded breeding owls in foothills





Summer 2015, 2016 and 2017

- Birds only in typical breeding areas
- No birds bred at wintering foothill sites;
All birds at higher elevations disappeared
- No wintering birds
from foothill sites at
typical breeding sites




Winter habitat differs from breeding season habitat

- Higher elevation
- Still need burrows, but simpler ones OK
- Fewer burrows, perhaps
- Single birds, perhaps
- Widely-distributed
- Low profile/cryptic



Results - 2014-2017



True Migrants! But, who are they?

Valuing and Protecting all Habitat



Birds pair up starting in February



Aggressive/Defensive

Typically seen when defending burrow



Females lay up to 12 eggs

Chicks stay below ground
for several weeks



Chicks emerge in May - stay with parents all summer




By September:

- * juveniles molt and disperse to seek their own burrows
- * adults typically migrate or move to other local burrows for the winter



Year-round Timeline



Burrowing Owls In Action!

- Adults at nest burrow:
- <http://www.arkive.org/burrowing-owl/athene-cunicularia/video-00.html>
- Parents and Chicks:
- <http://www.arkive.org/burrowing-owl/athene-cunicularia/video-03a.html>
- <http://www.arkive.org/burrowing-owl/athene-cunicularia/video-09.html>

Opportunistic predators

- * insects and small rodents dominate the diet
- * also eat amphibians, reptiles, crustaceans, birds



Diet in Santa Clara County, CA



Trulio, L. and P. Higgins. 2012. The diet of western burrowing owls in an urban landscape. *Western North American Naturalist* 72:348-356.


5 Study Sites: Total ~1450 ha (Site sizes: 62 to 722 ha)




Results—As a Whole

- 3092 pellets from 92 burrows
- 54 burrows associated w/specific owls

% Frequency - 94:6 (inverts to verts)



% Biomass: 30:70 (inverts to verts)



Key Findings

- Year-round prey - rodents and insects
- Composition and species, especially insect taxa, similar to other habitats
- CA vole and Botta's pocket gophers - do well in urban settings




Avg. mass = 53 g



Avg. mass = 155 g

Landscape as a Factor in Habitat Quality



Patches - yes - but enough foraging habitat in the landscape is needed to support long-term populations

Population Dynamics

- Adult survivorship: ~30-60% or more
- Juvenile survivorship: ~12-30%
- Nest success rates: Extremely variable
- Fecundity: Quite variable (~3 chicks per successful pair)
- PVA shows adult survivorship is the key parameter in population change (Barclay et al. 2011)

Population Genetics

- No genetic difference between migratory and resident birds
- Inbreeding due to isolated populations not evident
- Panmictic in the west
 - Migratory
 - Dispersal distances both short (1 mile or less) and long (50-150 miles or more)
(Results from Korfanta, et al. 2005)
- But...new data from fine-grained DNA tests

Small Group Exercise

Could they be here?

As a burrowing owl biologist, you are given information on a site. The owner wants to know, just based on these features, do you think there might be burrowing owls here?


Looking at these, what would you want to know about the site in order to say that owls might be there? What aspects of the site do you think would constrain or promote the presence of owls?

Status

- Endangered in Canada
- Threatened in Mexico
- Bird of Conservation Concern in US
- Endangered in Minnesota
- Threatened in Colorado
- Species of Special Concern in California, Montana, Oklahoma, Oregon, Utah, Washington, and Wyoming


Owls are declining in California

- 60% of breeding groups found in the 1980s disappeared by the 1990s
- A species of special concern in California



Example: Santa Clara County

Percent of 111 Sites Occupied in 1988 by Burrowing Owls Lost,
Reduced, or Extant over Time



Between 1988
and 2002, 66%
of locations lost



Only ~50 pairs
of birds remain
in all of Santa
Clara County

Santa Clara
Valley Habitat
Plan

Example: San Diego County Status

- Population Status - 1970s/1980s, ~250-300 pr;
2003, 25-30 pr
- Protection Efforts & Recovery Efforts

Lincer and Bloom, 2007



Threats

- #1 - Urbanization of grasslands. Urban sites are subject to disturbance, habitat loss, and poor habitat conditions.
 - Development
 - Auto strikes
 - Exterminating rodents
 - Secondary poisoning
 - More mesopredators & corvids
 - Weed abatement & Tall grass
 - Recreationists & Dogs
 - Surface/soil disturbance





Threats

- #1 - Urbanization of agricultural land. Loss of agricultural lands will impact burrowing owl populations.
 - ~90% of pairs found in agricultural landscapes
 - One of the only California raptors that does well in agricultural areas



Threats

■ #2 - Agricultural Practices

- Conversion to vineyards
- Lining irrigation ditches
- Discing to eliminate weeds
- Exterminating rodents
- Secondary poisoning

■ #3 - Solar/wind Farms

- Loss of ag lands
- Direct mortality




CA Burrowing Owl Distribution*

2006-07 estimate=
9,187 (SE=2,346) pairs statewide


Very similar to statewide
estimate of ~10 years before

* Wilkerson, R.L. and R. B. Siegel. 2010.
Assessing changes in the distribution and
abundance of burrowing owls in California,
1993-2007. *Bird Populations* 10:1-36.



Human Population Growth Expected:

- * Middle Central Valley
- * Southern Central Valley
- * Western Mohave
- * Imperial Valley



Regulatory Framework

- Federal: Migratory Bird Treaty Act - prohibits the "take" of any migratory bird or body parts, nests, eggs or products
- Federal: Fish and Wildlife Conservation Act - Bird of Conservation Concern
- Federal: Endangered Species Act, Section 10 - Habitat Conservation Plans
- State: California Fish and Wildlife Code Section 3503.5 - prohibits the taking, possession or destruction of birds of prey, their nests or eggs. For this reason, any impacts to burrowing owls during the breeding season (February 1 to August 31) are in violation of this code, unless approved by the CDFW

Regulatory Framework

- State: ESA - California Species of Special Concern
- State: Natural Community Conservation Planning Act (1991) - takes a broad-based ecosystem approach to planning for the protection and perpetuation of biological diversity
- State: California Environmental Quality Act (CEQA) - requires evaluation of project impacts to Species of Special Concern; requires a "mandatory finding of significance" if impacts to rare, threatened or endangered species are likely to occur
- State: CDFW Staff Report on Burrowing Owl Mitigation (2012) - guide for determining owl presence and avoiding impacts to owls and their habitat

Determining Presence/Absence

- Employ only qualified biologists (species-specific experience, education, & field training)
- Survey all suitable habitat areas an adequate time before disturbance (breeding or wintering)
- Observe at sunrise or sunset for at least 3 hr
- Observe at least 3 days
- Survey entire site on foot for burrows/birds
- If burrowing owls are found, contact California Department of Fish and Wildlife



Line Transect Surveys - Very effective for smaller areas

How to Manage Habitat to Preserve Burrowing Owls

- Habitat Features (owls present)
- Principles for Establishing Sites (owls not present)
- BUOW Relocation Review
- The Long View for California BUOWs


Habitat Features for Enhancing Areas for Owls (owls present)

- Shoreline Burrowing Owl Preservation Plan
- San Jose/Santa Clara Water Pollution Control Plant Interim Plan


Principle 1:

Develop a long-term plan that sets aside adequate areas for burrowing owl protection and management; exclude disturbance activities.






TOP VIEW



FRONT VIEW



1

- A - plastic irrigation valve box
- B - 4" diameter perforated corrugated plastic pipe
- C - 6" square hollow concrete block
- D - chain or plastic rope marking location of nesting chamber on ground surface
- E - 5' - 6' perch post (optional)

Principle 2: Enhance sites for nesting with artificial burrows.



Principle 3: Enhance the site for ground squirrels by bringing in mounds of dirt (don't use good soil!) and encourage healthy ground squirrel populations.



Principle 4: Keep grass short (<6 inches)
around nesting burrows and remove trees.



Principle 5: Enhance foraging opportunities by creating a structurally heterogeneous prey habitat; no pesticides or poisons.

<https://www.flickr.com/photos/123882326@N04/>



Recap - Key Habitat Features




- Open grassland habitat, few to no trees or other obvious raptor-perching sites
- As large as possible - viable site size will vary depending habitat quality and qualities of the surrounding landscape
- Healthy, breeding ground squirrel population
- Lots of burrows
- Short grass (<6") around burrows
- Structurally heterogeneous habitat—longer grass, foraging areas--for strong prey base

Management & Protection

Owls can do well in developed, urban, & agricultural areas if...

- Nests are protected from disturbance
- And there is enough foraging habitat

They don't need "pristine habitat"



Principles for Establishing Habitat - owls not present

- GOAL: Attract nesting owls to a site where they are not currently found
- NOTE: Once owls are extirpated from an area, reestablishing them is very difficult

Establishing Habitat

- Sites with the best chance of *attracting* nesting burrowing owls:
 - Add to adjacent, owl-occupied nesting habitat or within 300m of occupied habitat
 - Nesting owls recently on the site
 - Relatively large (~30-140 acres/owl pair??)
 - Not fragmented with roads or paths
 - Low elevation and flat
 - Habitat features as noted previously

Monitoring for Success

- Stable population over the years
- >50% of nests per year produce chicks
- Average of 3 chicks fledged per nest
- Some birds show site fidelity
- Acceptable levels of predation
- Successful habitat management for grass height and heterogeneity
- Strong prey base

Small Group Exercise

What are your recommendations for
habitat enhancing habitat for
burrowing owls?



What about relocating owls?

GOAL is to attract owls


Relocate birds only when absolutely necessary



Steve Blackmon, WildCare

Soft-release ("hacking") Set up

A.



B.



C.



Relocation Research Findings

- 105 wild, preflight juveniles soft-released at burrows in Minnesota 1986-1989 (Martell et al., 2001):
 - No birds ever found after release.
- 106 captive-raised, 10mo juveniles hard-released at burrows in British Columbia 1992-1997 (Leupin and Low, 2001):
 - 34% killed by predators
 - 2 overwintered for 3 years
 - 2 returned to release site after Spring migration
 - 7 successful nest attempts

Relocation Research Findings

- 27 adult birds moved from construction sites, soft-released at burrows in Santa Clara County in 1990s (Trulio, 1995):
 - 17 disappeared (63%) within a year of release
 - 7 birds (26%) flew back to their original site
 - 2 bred successfully on site (7%)
 - 1 victim of predation (4%)
- Researchers compare hard- vs. soft-release of captive-bred owls (2001-04) (Mitchell et al., 2011):
 - Soft-release results in greater survivorship and reproduction
 - 3% of adults returned the next year
 - 7% of chicks returned
 - 48% pairs fledged young; ~2.4 young/pair

Release conditions that seem to work best:

- Captive-reared, yearling adult owls
- One male and one female per burrow
- Birds reared in captivity near release sites
- Beginning of each breeding season
- Soft-release with birds in enclosures 14-17 days
- Supplemental feeding over the breeding season to maximize reproductive output.

The Long View for Burrowing Owls: Climate Change

How will the burrowing owl fare in an era of climate change?



Consider vegetation change*

* Lenihan, J.M., D. Bachellet, R.P. Neilson and R. Drapek. 2008. Response of vegetation distribution, ecosystem productivity, and fire to climate change scenarios for California. *Climate Change* 87 (Suppl. 1):S215-S230.

By 2100, under 3 climate change scenarios:


>70% increase in grasslands

Replacing shrub & Mixed evergreen woodland



Climate Change - Fire!

- 9-16% increase in area burned (Lenihan, et al., 2008)
- Fire frequency twice the current rate (Fried, J.S., M. Torn & E. Mills. 2004. *Climate Change* 64:169-191.)



Lenihan, et al. (2008)

Can Burrowing Owls Adapt?

- Live in range of habitats
- Increased grassland & fire could be exploited by burrowing owls
- But, climate change modeling suggests major losses to breeding habitat in the US.



Big Questions...



Dave Taylor, WildCare

- Fire + Habitat Change
 - Too much, too often?
 - Squeeze owls into less suitable conditions?
- And what about the intersection of human activities, population growth + climate change?
- National Audubon Climate Report states:


"By 2080, this diurnal owl species could lose 77 percent of its current breeding range. Climate change will disrupt its winter range as well, leaving only 33 percent intact..."

(<http://climate.audubon.org/birds/burowl/burrowing-owl>)

Our Challenge



+



Develop
Local Plans

Protect & Enhance
Current
Habitat

Identify
Future Owl
Habitat

Predict
Protect
Enhance

**Provide
conditions
for owls to
persist**

Thanks to colleagues & supporters

Especially...

- Phil Higgins, Debra Chromczak, Sandra Menzel
- Edmund Sullivan, Santa Clara Valley Habitat Agency
- City of Mountain View, Shoreline at Mountain View
- City of San Jose, WPCP
- NASA Ames Research - Moffett Federal Airfield
- US Fish & Wildlife Service - Don Edwards SFBNWR
- Santa Clara Valley Open Space Authority
- Santa Clara County Parks & Recreation, Santa Clara Valley Water District, and VTA

...and many tireless, enthusiastic field & lab assistants!

And thank you...

- Elkhorn Slough Coastal Training Program
- All the biologists, advocates, agency experts working to protect burrowing owls
- And, you for your attending this workshop to learn about this wonderful animal!



Photo by Ru